

strain of *P. luminescens* (W14) and then tested the effect of the mutant bacterial broths in our oral bioassay. Deletion of either *tca* or *tcd* individually (as *tca*⁻ or *tcd*⁻ mutant strains) greatly reduced the oral toxicity of the broth to *M. sexta*, whereas deletion of both *tca* and *tcd* together (in the *tca*⁻/*tcd*⁻ double mutant) eliminated oral toxicity altogether. These results suggest that both Tca and Tcd are involved in oral toxicity to Lepidoptera. However, we have been unable to purify sufficient quantities of Tcd to perform an LD₅₀ determination.

In order to examine the effects of Tca on the lepidopteran gut and compare it to that previously documented for both the *B. thuringiensis* δ -endotoxins and vegetative insecticidal proteins (Vips), and for cholesterol oxidase,^{11–15} we sectioned *M. sexta* neonates at intervals after oral ingestion of toxin. After several hours, toxin-treated midguts showed an accelerated rate of epithelial blebbing (Fig 2). This blebbing of the midgut epithelium into the lumen continues until the basement membrane is exposed and the epithelium is essentially destroyed. Both the columnar cells and the goblet cells appear to be attacked. Interestingly, a similar histopathology can be observed following injection of Tca directly into the insect haemocoel, which is presumably the normal route of delivery of the toxin by the bacterium.¹⁶

In conclusion, we have purified four toxin complexes from the culture broth of *P. luminescens* and cloned the four toxin complex-encoding loci. Genetic knockout of either *tca* or *tcd* reduces oral toxicity to *M. sexta* and knockout of both loci eliminates activity. Purified Tca shows effects specifically on the insect midgut, despite its putative normal delivery directly into the insect haemocoel. These *Photorhabdus* toxins (Phts) may form useful alternatives to other orally active bacterial protein toxins such as those from *B. thuringiensis* (Bt).

ACKNOWLEDGEMENTS

We thank all at Dow Agrosciences Biotechnology for their encouragement and support of this project. The work was supported by Hatch funds, The Applied Research and Technology Fund and The Industrial and Economic Development Fund, all administered by the University of Wisconsin-Madison and by Dow Agrosciences.

REFERENCES

- 1 Poinar GO, Thomas GM and Hess R, Characteristics of the specific bacterium associated with *Heterorhabditis bacteriophora* (Heterorhabditidae: Rhabditida). *Nematol* 23:97–102 (1977).
- 2 Dunphy GB and Webster JM, Virulence mechanisms of *Heterorhabditis heliothidis* and its bacterial associate, *Xenorhabdus luminescens*, in the non-immune larvae of the greater wax moth, *Galleria mellonella*. *Int J Parasitol* 18:729–737 (1988).
- 3 Gotz P, Borman A and Borman HG, Interactions between

- insects immunity and an insect-pathogenic nematode with symbiotic bacteria. *Proc R Soc Lond* 211B:330–350 (1981).
- 4 Schmidt TM, Bleakley B and Nealson KH, Characterization of an extracellular protease from the insect pathogen *Xenorhabdus luminescens*. *Appl Environ Microbiol* 54:2793–2797 (1988).
- 5 Dunphy GB and Webster JM, Lipopolysaccharides of *Xenorhabdus nematophilus* (Enterobacteriaceae) and their haemocyte toxicity in non-immune *Galleria mellonella* (Insecta: Lepidoptera) larvae. *J Gen Microbiol* 134:1017–1028 (1988).
- 6 Yamanaka S, Hagiwara A, Nishimura Y, Tanabe H and Ishibashi N, Biochemical and physiological characteristics of *Xenorhabdus* species, symbiotically associated with entomopathogenic nematodes including *Steinernema kushidai* and their pathogenicity against *Spodoptera litura* (Lepidoptera: Noctuidae). *Arch Microbiol* 158:387–393 (1992).
- 7 Clarke DJ and Dowds BCA, Virulence mechanisms of *Photorhabdus* sp strain K122 toward wax moth larvae, *J Invert Pathol* 66:149–155 (1995).
- 8 Bowen DJ and Ensign JC, Purification and characterization of a high molecular weight insecticidal protein complex produced by the entomopathogenic bacterium *Photorhabdus luminescens*. *Appl Environ Microbiol* (1998) in press.
- 9 Bowen D, Rocheleau TA, Blackburn M, Andreev O, Golubeva E, Bhartia R and French-Constant RH, Insecticidal toxins from the bacterium *Photorhabdus luminescens*. *Science (Washington)* 280:2129–2132 (1998).
- 10 Hofte H and Whitely HR, Insecticidal crystal protein of *Bacillus thuringiensis*. *Microbiol Rev* 53:242–255 (1989).
- 11 Sutter GR and Raun ES, Histopathology of European-Corn-Borer larvae treated with *Bacillus thuringiensis*. *J Invertebr Path* 9:90–103 (1967).
- 12 Kinsinger RA and McGaughey WH, Histopathological effects of *Bacillus thuringiensis* on larvae of the Indianmeal Moth and the Almond Moth. *Ann Entomol Soc Am* 72:787–790 (1979).
- 13 Endo Y and Nishiitsutsuji-Uwo J, Mode of action of *Bacillus thuringiensis* δ -endotoxin: histopathological changes in the silkworm midgut. *J Invertebr Path* 36:90–103 (1980).
- 14 Yu CG, Mullins MA, Warren GW, Koziel MG and Estruch JJ, The *Bacillus thuringiensis* vegetative insecticidal protein Vip3A lyses midgut epithelium cells of susceptible insects. *Appl Environ Microbiol* 63:532–536 (1997).
- 15 Purcell JP, Greenplate JT, Jennings MG, Ryerse JS, Pershing JC, Sims SR, Prinsen MJ, Corbin DR, Tran M, Sammons RD and Stonard RJ, Cholesterol oxidase: a potent insecticidal protein active against boll weevil larvae. *Biochem Biophys Res Comm* 196:1406–1413 (1993).
- 16 Blackburn M, Golubeva E, Bowen DJ and French-Constant RH, A novel insecticidal toxin from *Photorhabdus luminescens*: histopathological effects of toxin complex A (Tca) on the midgut of *Manduca sexta*. *Appl Envir Entomol* (1998) in press.

Synthesis of a biotin-like phosphonate model compound for (+)-hydantocidin

Werner Föry and Hans Tobler*
Novartis Crop Protection AG, Business Unit Herbicides,
R-1047.110, CH-4002 Basel, Switzerland

Abstract: Approaches to the synthesis of a biotin-like phosphonate are described. It was hoped that this would be a simpler model compound for the

* Correspondence to: Hans Tobler, Novartis Crop Protection AG, Business Unit Herbicides, R-1047.110, CH-4002 Basel, Switzerland
E-mail: hans.tobler@cp.novartis.com
(Received 1 July 1998; revised version 20 August 1998; accepted 16 February 1999)




Figure 1. Structures of (+)-hydantocidin (**1a**) and possible simpler model compounds.

naturally occurring spironucleoside (+)-hydantocidin, but it showed no activity as a herbicide nor as an inhibitor of AdSS.

Keywords: (+)-hydantocidin; model compound; phosphonate; biotin-like

1 INTRODUCTION

(+)-Hydantocidin (**1a**; Fig 1) is a naturally occurring spironucleoside with potent herbicidal activity and is produced by certain strains of the actinomycete *Streptomyces hygroscopicus* (Tü 2474).¹ It is actually a pro-herbicide, in that the corresponding 5'-phosphate (**1b**) inhibits adenylosuccinate synthetase (AdSS) in de-novo purine biosynthesis.^{1,2}

The presently known synthetic methods for the parent compound **1a** are not economically feasible and its close analogues lack biological activity.¹ Since X-ray data of the enzyme-inhibitor complex **1b** became available, the search for AdSS inhibitors has focused upon the CAMM-supported design of simpler model compounds: as an example, the biotin-like phosphonate **2** fulfils the steric requirements for an AdSS inhibitor.

Racemic **2** was synthesised in seven steps from the phosphonium salt **3a** (Fig 2), a close analogue of a recently identified biotin process key intermediate,³ by

using Wittig methodology for the introduction of the side chain (**5**) followed by appropriate functional group transformations and deprotection. In view of the well-known vigorous *N*-debenzylation reactions in biotin chemistry, a parallel synthesis using *para*-methoxybenzyl as a more-easily cleaved *N*-protecting group was carried out using **3b** as starting material. The choice of protecting group proved to be critical. The key intermediates **3** are readily produced in two steps from thiolactones **4**.³⁻⁷

2 SYNTHESSES

2.1 Introduction of side chain (Fig 3)

Wittig reaction of phosphoranes derived from *rac*-**3** with TBDMS-protected 2-hydroxy acetaldehyde^{8,9} competed with reductive formation of thiophanes **6** and **7**. The choice of base was crucial. Best results on a preparative scale (20 mmol) were achieved in the benzyl series with solid KN(TMS)₂ in the presence of a catalytic amount of 18-crown-6,¹⁰ thus affording the exocyclic allylic silyl ether **5a** in 66% yield (*E*-configuration predominant), accompanied by the thiophane **6a**.¹¹ On the other hand, an analogous Wittig reaction with **3b** under similar conditions proceeded disappointingly, **6b** and **7b** becoming the major components due to migration of the neighbouring *p*-MeO-phenyl moiety into the 2 α -position. Thus




Figure 2. Overall synthetic approach to model compound **2**.




Figure 3. Introduction of side-chain.




Figure 4. Side-chain transformations, I. (a): H₂, 10% Pd-C, MeOH, 10 bar, 50°, 18h; (b): TBAF, THF, 0°, 3.5h (c): CH₃P(OPh)₃⁺ I⁻, 2,6-Lutidin, DMF, 20°, 1.5h; (d): HPO(OEt)₂, KN(TMS)₂, THF, -70° to 20°, 20h; (e): P(Ph)₃, CCl₄, 20°, 20h; (f): P(OEt)₃, 130°, 16h, tlc-experiment.




Figure 5. Side-chain transformations, II. (a): Tetrabutylammoniumfluoride, THF, 0°, 3h; (b): P(Ph)₃, CCl₄, 75°, 20h; (c): P(OEt)₂(OTMS), 100°, 6h; (d): H₂, 10% Pd-C, MeOH, 10 bar, 50°, 18h.

the *p*-Meo-benzyl series was abandoned. Treatment of **3a** with base alone furnished **6a** as the sole product.

2.2 Side-chain transformations 1 (Fig 4)

Catalytic hydrogenation^{4,12} of olefin **5a** led to the mixture of epimers **8** and **12** in good yield, though with poor diastereoselectivity in favour of **12**. Chromatographic separation (elution with diethyl ether + hexanes, 1 + 1, by volume) and subsequent cleavage of the silyl ethers to alcohols **9** and **13**, posed no problems. The configuration at C-2 of either epimer was assigned on the basis of [¹H]NOE experiments.

Further treatment of **13** with methyltriphenoxyphosphonium iodide¹³ to obtain the corresponding iodide (as precursor for phosphonylation) afforded exclusively the hitherto-unknown tricycle **15** (a 5-thia-1, 2-diaza-cyclopenta[*cd*]pentalene system) as a viscous oil. With chloride **14** under Arbuzov conditions no phosphonate was observed but it again gave **15**. In contrast, the epimeric 2 β -alcohol **9** reacted readily to give the iodide **10**, from which the 2 β -phosphonate **11** was isolated and characterised.

The results in the 2 α -series may be explained in terms of neighbouring group participation. The close proximity of a benzyl group to the phosphonium intermediates (same type in both reactions) favours attack of halide ion at the benzylic carbon, resulting in ring closure and elimination of benzyl halides.

These facts suggest a reversal of the reaction order in

the side-chain transformations with phosphonylation at the allylic stage.

2.2 Side-chain transformations 2 (Fig 5)

Deprotection of silyl ether **5a** under standard conditions afforded the allylic alcohol **16** in good yield. Transformation to the chloride **17** proceeded cleanly but with reversal of configuration of the double bond (*Z*; NOE between olefinic *H* and *H*-3). Subsequent Arbuzov reaction gave the respective *Z*-configured allylic phosphonate **18**. In contrast to *E*-configured **5a**, catalytic hydrogenation^{4,12} of (*Z*)-phosphonate **18** proceeded with high diastereomeric excess, resulting in our proposed fully protected model 2 α -phosphonoethylthiophane **19**. Traces of 2 β -**19**, detected in [¹H]NMR, were identical with **11**.

2.3 Deprotection (Fig 6)

Deprotection of **19** proved to be a difficult task. After numerous failures, debenzylolation was achieved by simple treatment of **19** with concentrated sulfuric acid overnight at ambient temperature. Column chromatography on silica gel allowed purification and separation from regioselectively monodeprotected **20** to give the crystalline diethyl phosphonate **21** (mp 128–133 °C) in moderate yield. Finally, the pure racemic phosphonic acid **2** was obtained as a solid (mp 240–275 °C dec) after treatment with excess TMSBr and an aqueous work-up.




Figure 6. Deprotection. (a): conc.(96%) H₂SO₄, 20°, 16h; (b):(1) TMSBr (15 equiv.) DCM, 20°, 2.5 days, (2) H₂O.

3 BIOLOGICAL ACTIVITY

Tested at 250 µM **rac-2** and **rac-21** were inactive as inhibitors of adenosylsuccinate synthetase (AdSS).¹

Both compounds were also inactive when tested as herbicides against a selection of mono- and dicotyledonous weed species at 2 kg ha⁻¹ in the Novartis 1a standard screen.

REFERENCES

- 1 Fonné-Pfister R, Chemla P, Ward E Girardet M, Kreuz KE, Honzatko RB, Fromm HJ, Schä HP, Grütter MG and Cowan-Jacob SW, The mode of action and the structure of a herbicide in complex with its target: Binding of activated hydantocidin to the feedback regulation site of adenylosuccinate synthetase. *Proc Natl Acad Sci USA* **93**:9431–9436 (1996) and References 4–25 cited therein.
- 2 Heim DR, Cseke C, Gerwick BC, Murdoch MG and Green SB, Hydantocidin: A possible proherbicide inhibiting purine biosynthesis at the site of adenylosuccinate synthetase. *Pesticide Biochemistry and Physiology* **53**:138–145 (1995).
- 3 Eyer M (Lonza AG), Verfahren zur Herstellung von 1,3-substituierten Tetrahydro-1H-thieno-[3,4-d]-imidazol-2(3H)-on-4-yliden-pentansäureester. Eur Pat Appl 387 747 (1990).
- 4 Goldberg MW and Sternbach LH (Hoffmann-La Roche Inc), Synthesis of biotin. US Patent 2 489 232 (1949).
- 5 Goldberg MW and Sternbach LH, (Hoffmann-La Roche Inc), Imidazolidotetrahydrofurans. US Patent 2 489 233 (1949).
- 6 Gerecke M, Zimmermann J-P and Aschwanden W, Versuche zur Biotinsynthese. Herstellung von (3aS,6aR)-1,3-Dibenzyl-tetrahydro-4H-thieno[3,4-d]imidazol-2,4(1H)-dion. *Helv Chim Acta* **53**:991–999 (1970).
- 7 Seiter JJ, Process for producing 2-oxo-1,3-dibenzyl-4,5-cis-imidazolidinedicarboxylic acid. US Patent 5 151 525 (1992).
- 8 Sodeoka M, Yamada H and Shibasaki M, A new method for the stereocontrolled synthesis of silyl dienol ethers using (naphthalene)chromium tricarbonyl catalyzed isomerization. *J Amer Chem Soc* **112**:4906–4911 (1990).
- 9 Jung ME and Truc VC, Intramolecular Diels–Alder cycloadditions of substituted furfuryl E-2(phenylsulfonyl)acrylates. *Tetrahedron Lett* **29**:6059–6062 (1988).
- 10 Baker R and Sims RJ, A convenient modification of the Wadsworth–Emmons method of olefin formation using a crown ether. *Synthesis*, 117–119 (1981).
- 11 Lavielle S, Bory S, Moreau B, Luche ML and Marquet A, A total synthesis of biotin based on the stereoselective alkylation of sulfoxides. *J Amer Chem Soc* **100**:1558–1563 (1978).
- 12 Moolenaar MJ, Speckamp WN, Hiemstra H, Poetsch E and Casutt M, Synthese von D-(+)-Biotin durch selektiven Ringschluss von N-Acyliminium-Silylenolethern. *Angew Chem* **107**:2582–2584 (1995).
- 13 Verheyden JPH and Moffat JG, Halo sugar nucleosides. I. Iodination of the primary hydroxyl groups of nucleosides with methyltriphenoxyphosphonium iodide. *J Org Chem* **35**:2319–2326 (1970).

Plant cell cultures as a model for the biochemistry of crop selectivity

Peter Babczinski*

Bayer AG, Agricultural Centre Monheim, Institute for Metabolism Research and Residue Analysis, D-51368 Leverkusen, Germany

Abstract: Plant cell cultures have been used to study the metabolic degradation of 4-amino-5-methyl-2-(*tert*-butylaminocarbonyl)-1,2,4-triazolin-3-one. The biochemical basis of selectivity was shown to reside in effective metabolic conjugation. The herbicide was eliminated from cell cultures of beet (which is tolerant to it) by conjugation (N-glycolisation), but this occurred to only a limited extent with cell cultures of non-target plants such as soybean.

Keywords: crop selectivity; cell cultures; herbicide metabolism

Plant cell suspension cultures are well suited for the study of the metabolic degradation of xenobiotics.^{1–3} Here, studies on 4-amino-5-methyl-2-(*tert*-butylaminocarbonyl)-1,2,4-triazolin-3-one (**1**; see Fig 1), which is selective in sugar beet and shows broad-spectrum weed control against monocotyledonous and dicotyledonous species, using plant cell cultures, are presented. The aim of the studies was (i) to establish the pathway of metabolization in target (red beet) and non-target (soybean) plant cell suspension cultures, (ii) to investigate the biochemical basis for tolerance in the target cell culture and for selectivity between target and non-target cell cultures, and (iii) to compare the in-vitro results with the field situation.

Either dark-grown heterotrophic suspension cultures of soybean (*Glycine max* Merr cv *Merill v Mandarin*) or light-grown heterotrophic suspension cultures of red beet (*Beta vulgaris conditiva* Alef) were grown in Gamborg B5 medium containing sucrose, (20 g litre⁻¹), and 2,4-D, (0.1 mg litre⁻¹). The ¹⁴C-radiolabelled compound was applied in acetonitrile solution. Metabolites were isolated from either the nutrient medium or the cell extract (acetonitrile + water, 8 + 2 by volume), both after partitioning against

* Correspondence to: Peter Babczinski, Bayer AG, Agricultural Centre Monheim, Institute for Metabolism Research and Residue Analysis, D-51368 Leverkusen, Germany
(Received 29 June 1998; accepted 16 February 1999)